



Mental Health Review Journal

Trauma-informed mental healthcare in the UK: what is it and how can we further its development?

Angela Sweeney, Sarah Clement, Beth Filson, Angela Kennedy,

Article information:

To cite this document:

Angela Sweeney, Sarah Clement, Beth Filson, Angela Kennedy, (2016) "Trauma-informed mental healthcare in the UK: what is it and how can we further its development?", Mental Health Review Journal, Vol. 21 Issue: 3, pp.174-192, <https://doi.org/10.1108/MHRJ-01-2015-0006>

Permanent link to this document:

<https://doi.org/10.1108/MHRJ-01-2015-0006>

Downloaded on: 27 June 2018, At: 06:55 (PT)

References: this document contains references to 105 other documents.

The fulltext of this document has been downloaded 16520 times since 2016*

Users who downloaded this article also downloaded:

(2016), "Multidisciplinary team meetings in community mental health: a systematic review of their functions", Mental Health Review Journal, Vol. 21 Iss 2 pp. 119-140 <<https://doi.org/10.1108/MHRJ-03-2015-0010>><https://doi.org/10.1108/MHRJ-03-2015-0010>

(2012), "Despite the evidence – why are we still not creating more trauma informed mental health services?", Journal of Public Mental Health, Vol. 11 Iss 1 pp. 5-9 <<https://doi.org/10.1108/17465721211207266>><https://doi.org/10.1108/17465721211207266>

Access to this document was granted through an Emerald subscription provided by All users group

For Authors

If you would like to write for this, or any other Emerald publication, then please use our Emerald for Authors service information about how to choose which publication to write for and submission guidelines are available for all. Please visit www.emeraldinsight.com/authors for more information.

About Emerald www.emeraldinsight.com

Emerald is a global publisher linking research and practice to the benefit of society. The company manages a portfolio of more than 290 journals and over 2,350 books and book series volumes, as well as providing an extensive range of online products and additional customer resources and services.

Emerald is both COUNTER 4 and TRANSFER compliant. The organization is a partner of the Committee on Publication Ethics (COPE) and also works with Portico and the LOCKSS initiative for digital archive preservation.

*Related content and download information correct at time of download.

Trauma-informed mental healthcare in the UK: what is it and how can we further its development?

Angela Sweeney, Sarah Clement, Beth Filson and Angela Kennedy

The authors' affiliations can be found at the end of this article.

Abstract

Purpose – The purpose of this paper is to describe and explain trauma-informed approaches (TIAs) to mental health. It outlines evidence on the link between trauma and mental health, explains the principles of TIAs and their application in mental health and explores the extent to which TIAs are impacting in the UK.

Design/methodology/approach – The approach is a conceptual account of TIAs including a consideration of why they are important, what they are and how they can become more prevalent in the UK. This is supported by a narrative overview of literature on effectiveness and a scoping of the spread of TIAs in the UK.

Findings – There is strong and growing evidence of a link between trauma and mental health, as well as evidence that the current mental health system can retraumatise trauma survivors. There is also emerging evidence that trauma-informed systems are effective and can benefit staff and trauma survivors. Whilst TIAs are spreading beyond the USA where they developed, they have made little impact in the UK. The reasons for this are explored and ways of overcoming barriers to implementation discussed.

Originality/value – This paper – authored by trauma survivors and staff – describes an innovative approach to mental health service provision that, it is argued, could have immense benefits for staff and service users alike.

Keywords Mental health services, Childhood trauma, TIA, TIC, Trauma survivors, Trauma-informed

Paper type Conceptual paper

Received 30 January 2015

Revised 11 June 2015

7 April 2016

Accepted 1 June 2016

© Angela Sweeney, Sarah Clement, Beth Filson and Angela Kennedy.

Published by Emerald Group Publishing Limited. This article is published under the Creative Commons Attribution (CC BY 3.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licenses/by/3.0/legalcode>.

Angela Sweeney is funded by a National Institute for Health Research Post-Doctoral Fellowship. This paper presents independent research partially funded by the National Institute for Health Research (NIHR). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Introduction

It is known that many people in contact with mental health services have experienced physical or sexual trauma (Mauritz *et al.*, 2013), that there is a strong link between childhood trauma and adult mental distress (Bentall *et al.*, 2014), and that experiences of marginalisation, poverty, racism and violence are correlated with poor mental health (Paradies, 2006). This has led to a call for services to acknowledge psychological and social factors in the development of extreme mental distress (Read *et al.*, 2009). The hope is that such models would minimise the risk that people presenting to services have their symptoms disconnected from the context of their lives. In this paper, we will describe the concept of trauma-informed approaches (TIAs) which were developed in North America but have relatively few published models from public services across Europe. TIAs are based on the understanding that most people in contact with human services have experienced trauma, and this understanding needs to permeate service relationships and delivery (Harris and Falloot, 2001). We begin by examining the theoretical basis for TIAs including the link between trauma and mental distress and institutional retraumatisation. We will argue for a more systematic transformation of mental health services that acknowledges the role of trauma in people's lives and consequently reconceptualises relationships between survivors (people who have experienced trauma and mental distress and who may use mental health services) and

service providers. Finally, we present a narrative overview of literature on effectiveness of TIAs, map current TIA activity, explore why TIAs have not impacted on mainstream UK practice and discuss what might be needed to bring TIAs to the UK.

Defining trauma

Definitions of trauma vary, but broadly, trauma refers to events or circumstances that are experienced as harmful or life-threatening and that have lasting impacts on mental, physical, emotional and/or social well-being (SAMHSA, 2014). Trauma can be a single event or multiple events compounded over time. The concept of trauma encompasses experiences of interpersonal violence, such as rape or domestic violence. Complex childhood and developmental traumas include community violence (e.g. bullying, gang culture, sexual assault, homicide, war), abuse, neglect, abandonment and family separation (Van der Kolk, 2005; www.nctsn.org/trauma-types/complex-trauma). Lesser understood forms of trauma include social trauma, such as inequality, marginalisation, racism and poverty, and historical trauma, the trauma legacy of violence having been committed against entire groups, including slavery, genocide and the Holocaust (Blanch *et al.*, 2012). Lenore Terr (1991) has conceptualised two basic types of childhood trauma: Type I trauma involves witnessing or experiencing a single event such as a serious accident or rape. Type II trauma results from repeated exposure to extreme external events, such as ongoing sexual abuse.

Prevalence of trauma

The Adverse Childhood Experiences (ACE) study investigated the association between childhood trauma and adult health in over 17,000 people (predominantly white, middle class Americans; www.cdc.gov/violenceprevention/acestudy/prevalence.html). Childhood trauma was common: 30 per cent of respondents reported substance use in their household; 27 per cent reported physical abuse; 25 per cent reported sexual abuse; 13 per cent reported emotional abuse; 17 per cent reported emotional neglect; 9 per cent reported physical neglect; and 14 per cent reported seeing their mother treated violently (www.cdc.gov/violenceprevention/acestudy/prevalence.html).

Research has demonstrated that people in contact with the mental health system have experienced higher rates of interpersonal violence than the general population. A systematic review estimated that half of those in the mental health system had experienced physical abuse (range 25-72 per cent) and more than one-third had experienced sexual abuse (range 24-49 per cent) in childhood or adulthood, significantly higher than in the general population (Mauritz *et al.*, 2013). Similarly, survey research has found that people using mental health services are substantially more likely to have experienced domestic and sexual violence in the previous year compared to the general population (27 per cent of women and 13 per cent of men had experienced domestic violence compared to 9 and 5 per cent, respectively, of the general population; 10 per cent of women had experienced sexual violence compared to 2 per cent of the general population; Khalifeh *et al.*, 2015).

The link between trauma and mental health

Over the last decade, research evidence has increasingly supported the notion that trauma is linked to adult psychosis and a wide range of other forms of mental distress (e.g. Bentall *et al.*, 2014; Fisher *et al.*, 2010; Kessler *et al.*, 2010; Paradies, 2006; Varese *et al.*, 2012). The ACE study found that the more adverse life events people experience prior to the age of 18, the greater the impact on health and well-being over the lifespan, including poor mental health, severe physical health problems, sexual and reproductive health issues, engaging in health-risk activities and premature death (Anda *et al.*, 2010). Similarly, Shevlin *et al.* (2008) found that experiencing two or more trauma types significantly increased the likelihood of experiencing psychosis. Dillon *et al.* (2012) report evidence of a dose-dependent relationship between the severity, frequency and range of adverse experiences and subsequent impact on mental health. Interestingly, research has also demonstrated that the general public share the notion that trauma and adverse life events play a causal role in mental health difficulties (e.g. Read *et al.*, 2013; Angermeyer and Dietrich, 2006).

Contemporary neuroscience is exploring the link between childhood trauma and neurological development. This research is informing TIAs which typically adopt a whole systems view of people and their environments, including an understanding of the role and impact of neurological damage. For instance, research has demonstrated that trauma has an impact on developing brains in childhood which can go on to affect the structure and function of adult brains (Perry 1995, 2005). This has led to the development of a traumagenic neurodevelopmental understanding of the link between childhood adversity and adult psychosis, which now has a large body of supporting evidence (Read *et al.*, 2014). The neurological damage caused by trauma suggests that survivors can be “primed” to respond to current situations that replicate the experience of loss of power, choice, control and safety in ways that may appear extreme, or even abnormal, when a history of past adverse events is not taken into account. However, research has also indicated the healing potential of current relationships (Perry, 2005). To find out more see, for example, Van der Kolk (2005), Read *et al.* (2014) and Dillon *et al.* (2012).

Research has also demonstrated that traumatic events are more frequently experienced by people in low-socioeconomic groups and from minority ethnic communities (e.g. Hatch and Dohrenwend, 2007). It has further been argued that poverty is the most powerful predictor of mental distress because it predicts so many other causes (Read, 2010). Moreover, black people are over-represented in the mental health system, are more likely to experience negative or adversarial pathways to care, to be diagnosed with psychotic disorders and to receive compulsory treatment (e.g. Mohan *et al.*, 2006; Morgan *et al.*, 2004). Yet, there is little discussion of the potential role of historical and cultural trauma in this. Indeed, social trauma, including poverty, racism and urbanicity, is so prevalent it is often not recognised as integral to poor mental health by clinicians or those experiencing it.

Notably, people in contact with mental health services who have been sexually or physically abused in childhood typically have longer and more frequent hospital admissions, are prescribed more medication, are more likely to self-harm and are more likely to attempt to kill themselves than people without experiences of childhood abuse (Read *et al.*, 2007).

Retraumatization in the mental health system

Retraumatization essentially means to be traumatised again. It occurs when a person experiences something in the present that is reminiscent of a past traumatic event. This current event or trigger often evokes the same emotional and physiological responses associated with the original event. People are not always aware that their current distress is rooted in past events, nor do all people relive the original event in a logical, coherent manner (Durant, 2011).

The mental health system can retraumatise survivors through its fundamental operating principles of coercion and control (Bloom and Farragher, 2010). Retraumatization includes overt acts, such as restraining and forcibly medicating a rape victim, as well as less palpable retraumatization, such as pressure to accept medication which mimics prior experiences of powerlessness. Empirical research indicates that traumatic experiences (e.g. physical assault, seclusion, restraint) are widespread in inpatient settings (Freuh *et al.*, 2005). Mental health services can also contribute to historical and cultural trauma by recasting responses to racism as individual pathology (Jackson, 2003), recasting women’s attempts to resist domestic control as hysteria (St-Amand and LeBlanc, 2013) and recasting homosexuality as sexual deviance in need of corrective treatment (Friedman, 2014).

Jennings believes that whilst retraumatization can be unintentional and unanticipated, it will remain whilst mental health systems fails to acknowledge the role of trauma in people’s lives and their consequent need for safety, mutuality, collaboration and empowerment (www.theannainstitute.com). Current services and supports that do not take these impacts into account may inadvertently retraumatise, further reinforcing survivors’ needs for coping strategies such as illicit drug use or self-harm.

The impact of retraumatizing systems on staff

The policies, procedures and practices that staff may be required to perform in “trauma-organised systems” (Bloom and Farragher, 2010) can conflict with personal and ethical codes of conduct. For example, the use of seclusion and restraint as an institutional practice erodes the

very meaning of compassion and care, the primary reasons most staff enter their chosen field. Staff who experience conflicts between job duties and their moral code are under chronic stress for which they must learn to cope and adapt. Those coping strategies may include “shutting off” the ability to empathise, and viewing people receiving services as “other” thereby disqualifying their humanity and basic human rights. Pessimism – rather than enthusiasm and hope – may buffer staff from their own feelings of helplessness (Chambers *et al.*, 2014).

Staff may also engage in “power over” relationships when organisations place a higher priority on risk management than human relationships. A nurse who is required to perform a personal search may become frustrated by a service user’s resistance, failing to recognise that she/he is a stranger who is placing hands on the body of another who may be a rape survivor. Organisational cultures may become corrupted, paving the way to power over relationships that reinforce people’s helplessness and hopelessness. In these “corrupted cultures”, the basic values of the organisation are no longer driving practice; instead, the needs of service users become secondary to the needs of staff, and restraint and coercion may be used widely even when less restrictive options are available. This and other working practices and routines (such as rigid professional hierarchies and a lack of supervision) can dehumanise both staff and service users and lead to human rights violations (for an account of corrupted cultures and the impact on coercion see, Paterson *et al.*, 2013; Wardhaugh and Wilding, 1993). The National Institute for Clinical Excellence (NICE) has expressed frustration at first resort to coercive practices even where other approaches are indicated (NICE, 2005). The impact of trauma-organised services on workers is analogous to the impact of trauma on survivors – it reshapes and re-constructs self-identity and can shatter individual meaning and purpose (Knight, 2015).

The principles of TIAs

The development of TIA can be traced to the USA and to Harris and Fallot’s (2001) seminal text, *Using Trauma Theory to Design Service Systems*. Bloom (2013), also from the USA, who developed the Sanctuary Model outlines the development of TIA from the era of moral treatment, through social psychiatry and finally the concept of the Therapeutic Community (Bloom and Norton, 2004) which includes developments in the UK. TIAs can be defined as “a system development model that is grounded in and directed by a complete understanding of how trauma exposure affects service user’s neurological, biological, psychological and social development” (Paterson, 2014). Consequently, TIAs are informed by neuroscience, psychology and social science as well as attachment and trauma theories, and give central prominence to the complex and pervasive impact trauma has on a person’s worldview and interrelationships.

TIAs are applicable to all human services, including physical health, education and schools, forensic, housing and social care (Schachter *et al.*, 2008; Havig, 2008; Cole *et al.*, 2013). In a trauma-informed service, it is assumed that people have experienced trauma and may consequently find it difficult to develop trusting relationships with providers and feel safe within services. Accordingly, services are structured, organised and delivered in ways that engender safety and trust and do not retraumatise. Thus, trauma-informed services can be distinguished from trauma-specific services which aim to treat the impacts of trauma using specific therapies and other approaches. The key principles underlying TIAs can be found in Table I, adapted from SAMHSA (2014), Elliot *et al.* (2005) and Bloom (2006).

Whilst it may seem that principles such as safety and collaboration define any good service for any service user, Elliot *et al.* (2005) have argued that if these principles are not adhered to, trauma survivors may be unable to use services. It is striking that these general principles have strong resonance with the values that psychiatric survivors have historically called for, and underpin much peer support practice (e.g. Mead and MacNeil, 2006).

What are the potential benefits of TIAs?

The potential benefits of TIAs to survivors are myriad, including hope, empowerment, support that does not retraumatise and access to trauma-specific services. Moreover, the medicalisation of human suffering has created a divide between people receiving services and those offering

Table 1 The key principles of trauma-informed approaches

1. Recognition	Recognise the prevalence, signs and impacts of trauma. This is sometimes referred to as having a trauma lens. This should include routine enquiry about trauma, sensitively asked and appropriately timed. For individual survivors, recognition can create feelings of validation, safety and hope
2. Resist retraumatisation	Understand that operational practices, power differentials between staff and survivors, and many other features of psychiatric care can retraumatise survivors (and staff). Take steps to eliminate retraumatisation
3. Cultural, historical and gender contexts	Acknowledge community-specific trauma and its impacts. Ensure services are culturally and gender appropriate. Recognise the impact of intersectionalities, and the healing potential of communities and relationships
4. Trustworthiness and transparency	Services should ensure decisions taken (organisational and individual) are open and transparent, with the aim of building trust. This is essential to building relationships with trauma survivors who may have experienced secrecy and betrayal
5. Collaboration and mutuality	Understand the inherent power imbalance between staff and survivors, and ensure that relationships are based on mutuality, respect, trust, connection and hope. These are critical because abuse of power is typically at the heart of trauma experiences, often leading to feelings of disconnection and hopelessness, and because it is through relationships that healing can occur
6. Empowerment, choice and control	Adopt strengths based approaches, with survivors supported to take control of their lives and develop self-advocacy. This is vital as trauma experiences are often characterised by a lack of control with long-term feelings of disempowerment
7. Safety	Trauma engenders feelings of danger. Give priority to ensuring that everyone within a service feels, and is, emotionally and physically safe. This includes the feelings of safety engendered through choice and control, and cultural and gender awareness. Environments must be physically, psychologically, socially, morally and culturally safe
8. Survivor partnerships	Understand that peer support and the coproduction of services are integral to trauma-informed organisations. This is because the relationships involved in peer support and coproduction are based on mutuality and collaboration
9. Pathways to trauma-specific care	Survivors should be supported to access appropriate trauma-specific care, where this is desired. Such services should be provided by mental health services and be well resourced

support; this divide can create tenuous bonds that are inadequate, at times, to protect the human and civil rights of people viewed as other (Filson and Mead, 2016). But trauma is something that many of us experience, and indeed, a small number of studies suggest that workers in human services have a high prevalence of ACE scores (e.g. Esaki and Larkin, 2013). In recognising trauma as a shared event, healing too becomes something we do together.

Because TIAs are premised on the understanding that most of the people who come into contact with mental health services have been impacted by trauma, training, supervision and support for staff are seen as essential. This attention to staff support has the potential to decrease burnout and reduce staff turnover. For example, research suggests that supervisors who feel that their organisation values them and cares about their well-being are more likely to be supportive towards the people they are responsible for (Shanock and Eisenberger, 2006).

There are complex interactions between service users, practitioners and organisations that can come to mirror one another through “parallel processes” (Bloom, 2006). Trauma survivors’ lives may be organised around the trauma experience, just as systems can come to be organised around models that are inadequate for responding to survivors. This means that, for example, in trauma-organised systems, survivors may feel and be unsafe, leading to aggression towards staff. Experiencing aggression from survivors may cause staff to become wary and hostile, with organisations responding with greater punitive and risk-averse measures. This increases survivors’ sense of unsafety and aggression. Becoming trauma-informed has the potential to break these negative parallel processes and create positive interactions.

Trauma carries a heavy economic cost. Dolezal *et al.* (2009) have reviewed US research evidence on the economic impacts of violence and abuse and estimate a cost of between 17 and 37.5 per cent of the total spend on healthcare. They believe that a compassionate healthcare system that understands the impacts of violence and abuse and offers appropriate support may avoid many of these costs. In the UK, the Department of Health (DH) has estimated that:

Costs include the costs of providing public services for victims, the lost economic output of women and the human and emotional costs of violence for victims. An indicative figure for the minimum cost of violence against women and children is £36.7bn (DH, 2011).

There is also some evidence that a reduction in seclusion and restraint has large cost savings (e.g. a 92 per cent reduction in the costs linked to restraint, LeBel and Goldstein, 2006).

Applying trauma-informed principles to mental health

Trauma-informed mental health services are strengths based: they reframe complex behaviour in terms of its function in helping survival and as a response to situational or relational triggers. Reframing refers to looking at, presenting, and thinking about a phenomenon in a new and different way, and replaces traditional individual/medical model approaches to madness and distress with a social perspective, somewhat akin to the Social Model of Disability (Wilson and Beresford, 2002). Reframing behaviour as meaningful allows providers to address underlying needs and utilise less intrusive strategies. We have fictionalised a trauma-informed response to a woman who self-harms in Box 1.

In a trauma-informed mental health service, all staff – clinical and non-clinical – understand the impact of trauma on a person’s ability to survive in the present moment. Crucially, this entails a shift from thinking “what is wrong with you” to “what happened to you” (Harris and Fallot, 2001). The critical roles of racism, sexism, homophobia, ageism, poverty and their intersectionalities are recognised. Survivors in crisis are not viewed as manipulative, attention-seeking or destructive, but as trying to cope in the present moment using any available resource.

Providers do not fear asking about trauma, yet do so in ways that are respectful of potential retraumatisation; the power of telling one’s story but also the impotency of telling it where nothing changes (Filson, 2011); the need to move at the survivor’s pace; the need to truly listen; and the need for post-disclosure support. Survivors are forewarned about trauma questions, and can choose not to answer. Trauma information is integrated into treatment plans so that people can be referred to trauma-specific services (if wanted) (see Read *et al.*, 2007 for a full account of why, when and how to ask about abuse).

The basic safety of environments is prioritised – physical, psychological, social and moral – with organisations making a commitment to non-violence (Bloom, 2006). Staff receive support to help them focus on trauma, and steps are taken to build a sense of community and shared responsibility between staff and survivors (Bloom, 2006). This means that services prioritise building trusting, mutual relationships between staff and survivors. When relationships are prioritised, policies and procedures (such as time limited sessions with a therapist) can be re-evaluated in light of whether or not they support TIAs.

TIAs in mental health aim to reduce or eradicate coercion and control, including medication as restraint, verbal coercion, threats of enforced detention, withholding information, restrictive risk-averse practices, disrespectful and infantilising interactions and Community Treatment Orders (see, for instance O’Hagan, 2003). Clinicians understand the revictimisation that “power over” relationships reinforce. Training and supervision provide staff with the tools to attend to potential relational and situational triggers and to use trust-based, collaborative relationships to support people.

Box 1. Jenny

Jenny has had numerous hospital admissions over four years, usually through self-harming events, including swallowing foreign objects and cutting her arms. Previously, some staff described Jenny as “attention-seeking” and “manipulative”, and responded by trying to control or stop the behaviour. This included ignoring Jenny, giving PRN medication or forcibly medicating her. This has changed since the organisation began training its staff on trauma and trauma-informed approaches. Now, when Jenny tells staff she wants to hurt herself, staff respond to Jenny’s pain, recognising that past strategies added to Jenny’s sense of powerlessness. While Jenny’s safety is no less important, validating Jenny’s pain and her attempt to cope with it, along with using harm reduction strategies around her self-injury, has greatly helped her. Jenny is now using art to bring voice to her experiences, and her treatment team have referred Jenny to a therapist who will work with her on the issues arising from her experiences of abuse.

Survivors often encounter numerous human services across their lives. To be trauma-informed, each service within and beyond the local mental health system should operate according to TIA principles. This includes primary care, A&E, talking therapies, mental health teams, crisis care, the police, social services and voluntary sector services (such as trauma-specific service providers).

What is the evidence on the effectiveness of TIAs?

To provide an overview of the current state of evidence on the effectiveness of TIAs in mental health we searched nine electronic databases (Medline, Embase, PsycInfo, CINAHL, Cochrane Library, Sociological Abstracts, Social Policy and Practice, Global Health and Maternity and Infant Care) using the title-word search "trauma and informed". Searches were from the earliest date of each database to August 2014 (searches run on 5 August 2014). This yielded 129 unique publications. One author (SC) assessed the items against the following inclusion criteria: participants – adults and/or youth in receipt of mental health services; intervention – any form of trauma-informed care; control (any or none); outcomes – any, including those relating to staff as well as service users; and study design – any design providing evidence on effectiveness, including qualitative evaluation studies.

Eight studies (Azeem *et al.*, 2011; Chandler, 2008; Domino *et al.*, 2006; Gatz *et al.*, 2007; Greenwald *et al.*, 2012; Messina *et al.*, 2014; Morrissey *et al.*, 2005; Weissbecker and Clark, 2007) met the inclusion criteria. The findings are presented in Table II.

All studies were conducted in the USA, and four were evaluations of Women and Co-Occurring Disorders services. Four studies were controlled pre-post-studies, two were pre-post-studies and one was a qualitative study. Beneficial effects noted in these studies included reduction in seclusion, reduced post-traumatic stress symptoms and general mental health symptoms, increased coping skills, improved physical health, greater treatment retention and shorter inpatient stays (see Table II, main findings). Other outcomes did not change such as substance misuse, emergency room use, imprisonment and shelter use.

Several of the studies were large, multi-site and quasi-experimental. The evidence-base is limited, however, by the relatively small number of studies, restricted to one country, with half the studies evaluating one particular TIA model. We did not locate any randomised controlled trials of trauma-informed mental healthcare. The findings cannot necessarily be generalised beyond the USA where the evidence is located. Also, as noted by Greenwald *et al.* (2012), the inclusion of numerous interventions makes it difficult to precisely identify the causes of the improvement. Integrating trauma-specific services within a broader trauma-informed service is advocated from a theoretical and service perspective; therefore, as is the case with complex interventions, research may not be able to pinpoint the key active ingredients.

This narrative overview of the evidence has limitations. It is not a systematic review as to conduct one was beyond the scope and intention of this paper. Consequently, although the number of bibliographic databases searched was large, the search strategy was basic, we did not search the grey literature or use reference checking or consultation with experts, and the selection of studies was undertaken by one investigator. A future systematic review on the effectiveness of trauma-informed care generally or in a mental health context would be welcome and illuminating.

Future research will hopefully provide a fuller picture of what TIAs are able to achieve. Given the sound theoretical and ethical underpinnings of TIAs and the extensive developmental work undertaken, coupled with the current promising evidence to date, there is certainly a strong case for the wider implementation and evaluation of TIAs.

TIAs in the UK

The concept of TIAs originated in the USA and in 2005 the United States Federal Substance Abuse and Mental Health Services Administration (SAMHSA) established a National Centre for Trauma-Informed Care. Many human service providers, including those in mental health, are now familiar with the concept of TIAs. However, the US mental health system remains heavily biomedical and despite conceptual familiarity, implementation of TIAs across sectors is patchy

Table II Narrative review of evidence on trauma-informed mental health care: study characteristics and findings

Study	Study design	Participants and setting	Intervention	Main findings
Azeem <i>et al.</i> (2011)	Pre-post-study	458 youth admitted to a child and adolescent psychiatric hospital, USA	Training staff in six core strategies that are based on trauma-informed care	In the first six months of study, the number of seclusions/restraints episodes were 93 (73 seclusions/20 restraints), involving 22 children and adolescents. In final six months of study following the training programme, there were 31 episodes (six seclusions/25 restraints) involving 11 children and adolescents
Chandler (2008)	Qualitative interview study, content analysis	Ten staff in an inpatient mental health unit, USA	Introduction of a trauma-informed approach	Themes identified included: changing perspective, developing collaborative relationships, implementing safety measures and prescribing educational resources, which the authors concluded indicated significant cultural change
Domino <i>et al.</i> (2006)	Controlled pre-post-study	1,023 women with co-occurring mental health and substance abuse problems with histories of interpersonal violence, USA	Trauma-informed outpatient group counselling, in the context of integrated substance abuse and mental health services	Intervention group women used the internal services provided. The intervention did not have strong effects on patterns of service use external to the intervention, such as emergency room, jail and shelter use
Gatz <i>et al.</i> (2007)	Controlled pre-post-study	313 women with co-occurring mental health and substance use disorders and histories of trauma, USA	Seeking safety, a trauma-specific group treatment focusing on safety and coping skills, in the context of integrated substance abuse and mental health services	Intervention women showed significantly better treatment retention over three months and greater improvement on post-traumatic stress symptoms and coping skills
Greenwald <i>et al.</i> (2012)	Pre-post-study	53 youth in a residential treatment facility, USA	Training in the Fairy Tale model of trauma-informed treatment	Compared to the year prior to training, in the year of the training the average improvement in presenting problems was increased by 34 per cent, time to discharge was reduced by 39 per cent, and rate of discharge to lower level of care was doubled
Messina <i>et al.</i> (2014)	Controlled pre-post-study	277 women offenders, USA	Gender-responsive treatment	Significant reduction in PTSD
Morrissey <i>et al.</i> (2005)	Controlled pre-post-study	3,034 women with co-occurring mental health and substance use disorders in nine outpatient sites, USA	Comprehensive, integrated, trauma-informed and consumer-involved approach to treatment	For substance use outcomes, no effect was found. The meta-analysis demonstrated small but statistically significant overall improvement in women's trauma and mental health symptoms in the intervention relative to the usual-care comparison condition
Weissbecker and Clark (2007)	Controlled pre-post-study	2,189 women with co-occurring disorders and histories of violence	Comprehensive, integrated, trauma-informed and consumer-involved approach to treatment	Improved physical health outcomes

(Cheryl Sharp, Senior Advisor for Trauma-Informed Services, National Council for Behavioural Health, USA, personal communication). In terms of strategies and implementation, the International Initiative for Mental Health Leadership has produced a brief report on key national and regional activities in TIAs across nations involved with the IIMHL as part of their "Make it so" series (IIMHL, 2012). As the aim of this series is to rapidly share information, it is likely that some key activities have been missed. Notwithstanding this caveat, countries where trauma-informed practices were identified were the USA, Canada, Australia and New Zealand. Nascent strategies and practices were described for Scotland and Ireland with nothing noted for England. The USA was described as leading the world in TIAs, and as the only nation to have national policy relating to trauma. Specifically, in 2011-2014 the eight strategic objectives of the SAMHSA included

“trauma and justice”. This objective aimed to reduce the impacts of violence and trauma by integrating TIAs throughout health, behavioural health and related systems.

Of the 129 academic publications with “trauma” and “informed” in the title found when we conducted our overview of the evidence of effectiveness, the vast majority of publications (86 per cent) were from the USA. Five other countries, including the UK, had a small amount of academic activity. The UK work comprised two discussion papers (Rose *et al.*, 2012; Ardino, 2014), a book (Taylor, 2012) and two reviews (Harragan, 2013; Steckley, 2013) of a non-UK-authored book on TIA. The focus was typically on residential treatment or mental health services for children and adolescents, with two general mental health papers (Ardino, 2014; Rose *et al.*, 2012).

Academic publication rates may have little bearing on the provision of services, therefore to gain a wider perspective on the current UK situation, we conducted a Google search on UK pdf documents with “trauma-informed” in the title. This yielded information about:

- a trauma-informed foster care service in North Wales, Chester and the Wirral (www.newfocas.co.uk/includes/spaw2/uploads/files/trauma.pdf);
- a guide to trauma-informed resettlement for people leaving youth custody (www.beyondyouthcustody.net/wp-content/uploads/BYC-Developing-trauma-resettlement-youth-custody-practitioners-guide.pdf);
- workshops led by visiting North American experts (<http://stephaniecovington.com/assets/files/Brochure-Flyer-dated-2-26-14-Woman-at-Risk-Flyer-V1.pdf>; www.vptorg.co.uk/whoslistening/docs/conf%20speeches/Aimee_Neri_Presentation.pdf; www.stir.ac.uk/trauma14/);
- conferences, including two on mental health (www.makingwaves.org/news/trauma-informed-services-conference/; www.stir.ac.uk/trauma14/; www.bild.org.uk/our-services/events/supportingpeople/); and
- training with a trauma-informed approach (www.ukpts.co.uk/site/assets/CALM-Final.pdf; www.calmtraining.co.uk/index.php; www.ewrasac.org.uk/Training/; www.avaproject.org.uk).

One of us, (AK), has played a key role in introducing TIAs to Tees, Esk and Wear Valleys NHS Foundation Trust, and describes her experiences in Box 2.

It is clear that TIAs are beginning to reach the UK, although often in settings beyond mental health. However, the two conferences on trauma-informed mental healthcare in 2014 – with speakers from psychology, mental health nursing, psychiatry and the survivor movement – indicate the beginnings of a sea-change. Scotland’s Mental Health Strategy 2012-2015 includes psychological trauma as a key priority (Scottish Government, 2012). The strategy states that “General Services should be Trauma Aware”, and aims to improve recognition and awareness of trauma in Primary Care and Mental Health Services, encourage staff to make appropriate referrals for trauma survivors, and roll out trauma training. Although TIAs are not named, this is nevertheless a welcome development.

Similarly, the National Mental Health Development Unit (2010) and DH (2011) have released strategy documents on gender sensitive services that include trauma awareness. The DH published recommendations regarding routine enquiry of abuse in mental health settings over a decade ago (DH, 2003) and a programme of work was undertaken to train staff, which demonstrated changes in skill (McNeish and Scott, 2008). This focused on changing the emphasis from “What is wrong with this person?” to “What has happened to this person?”. Asking the basic question: “Have you ever experienced physical, sexual or emotional abuse at any time in your life?” has now become mandatory for UK services. However, current evidence that staff do this in practice is scant and this suggests that good practice that goes beyond this question is not widespread (Hepworth and McGowan, 2013; Brooker *et al.*, 2016). One significant change that may prompt responses from services is the inclusion of trauma in some NICE guidelines, for example, the recently updated guidance for the management of schizophrenia (NICE, 2014). Some early intervention services for psychosis, in particular, are attempting to be more trauma-informed. Toner *et al.* (2013) showed that having a formulation-driven approach to understanding psychosis was more important in creating

Box 2. Case study: introducing TIA to an NHS Trust

Tees, Esk and Wear Valleys NHS Foundation Trust (TEWW) is a large mental health provider in the North of England which serves a population of 1.6 million people and employs over 5,000 staff. TEWW is embarking on a program to develop trauma-informed services throughout its adult division. Its TIA has been to develop a pathway of care and to train staff to implement this pathway. Training is undertaken as a team and it has been well received with staff reporting it relevant to their work and increasing their confidence afterwards. The resource set incorporates a number of elements: it has a variety of information leaflets for clients; resource links and summaries for staff; a treatment algorithm; service skills matrix; good practice guidance for managing trauma disclosures; information on screening for dissociation and how to manage it; a section on staff well-being; and a framework for understanding risk issues. The pilot project on an acute adult mental health ward included all staff from senior medics to health care assistants. They found that three quarters of the people admitted could directly link trauma with their current difficulties. Totally, 80 per cent had substance misuse issues, and the same proportion self-harmed. Totally, 40 per cent were experiencing some psychosis. Ward staff felt empowered to have meaningful discussions about trauma and used this to inform formulation based care plans. They were able to implement some core skills in grounding and emotion regulation, which resulted in a reduction in the use of PRN medication. It was important that local trauma champions in each team facilitated supervision, management and implementation of the guidance. Staff could also call on external complex case consultation for trauma, which was evaluated as being extremely helpful. Follow-up training plans were then developed to respond to specific areas of need as requested and so far this has been dominated by dissociation. TEWW has promoted experts by experience to deliver much of this. Finally, trauma specific supervision groups are supporting therapists to respond to issues of complex trauma.

staff that were empowered to address trauma than having the ability to enquire about it. There is something very important about the model of mental health that staff bring with them to the role.

What are the barriers to implementing TIAs in the UK?

We have identified a number of potential explanations for the slow implementation of TIAs in the UK, although our list is not exhaustive. Many of these implementation barriers are applicable to settings beyond the UK. First, despite compelling evidence, there remains strong resistance to the notion that trauma and childhood abuse plays a causal role in psychosis and mental distress. Historically, such claims have been seen as “family blaming”, and have been vehemently opposed, e.g. historic opposition to Freud and Laing. Instead there is a focus on the biological basis of mental distress, with genes and neurology seen as causal and trauma relegated to a trigger at best (Moskowitz, 2011). Thus, mental distress is understood as a scientific, medical and pharmacological problem, rather than a human, familial or social issue.

Second, Western societies have strongly resisted notions of historical and cultural violence and their consequent trauma legacies. Jackson (2003), an African American survivor and therapist, has produced a powerful research account of scientific racism, slavery and colonialism and the impact this has had on survivors generationally and today. Focussing on the social and systemic causes of trauma places practitioners in opposition to powerful groups and consequently is often avoided (Coles, 2014).

Third, Coles (2014) has described “horror” as a barrier to practitioners embracing notions of trauma: “to stand as witness to the extent and horror of people’s accounts of pain and suffering is to encounter and experience fear, despair, loss and rage”.

Fourth, UK public services face continuous change and upheaval, making many wary and weary of new initiatives. Consequently, introducing new conceptualisations of care can be challenging, and this is particularly acute with TIAs because the role and prevalence of trauma is disputed (e.g. the DH and NICE focus on diagnostic categories, rarely referring to trauma). Compounding this, UK austerity means that resources are scarcer and morale lower. This context makes it harder to engage with new initiatives.

Fifth, TIAs are a relatively complex and involved approach to service provision, and are easily confused with trauma-specific services. Muskett (2014) has described how mental health nurses in Australia struggle to translate TIA principles into their everyday practice beyond reducing seclusion and restraint.

Sixth, there have been a number of initiatives aimed at improving mental health services and relationships between service providers and users. For example, in the UK, Star Wards aims to support excellence on inpatient psychiatric wards (www.starwards.org.uk), Safewards aims to reduce conflict and containment and increase safety on inpatient mental health wards (www.safewards.net/), and Compassion in Practice centralises the six Cs of nursing and midwifery (care, compassion, competence, communication, courage and commitment). Whilst such initiatives are compatible with TIA, they are nonetheless another way to conceptualise and implement care for providers to grapple with.

Seventh, many UK mental health staff have no access to regular structured supervision, and this is a serious barrier to the implementation of TIA. In our case study (Box 3) we cite trauma-specific supervision groups as a way of supporting therapists to respond to issues of complex trauma.

Finally, once a concept starts to take hold it can gain momentum. Debate, training opportunities, champions, mentors and networking all perpetuate thinking and practice. Our mapping work suggests that despite evidence of increasing interest in TIAs in the UK, we have not yet achieved the critical mass needed for frontline TIA implementation.

How can we bring TIAs to the UK?

Addressing each of the issues outlined above will go some way towards contributing to the development of UK TIA practice. First and foremost, a paradigm shift in collective thinking about

Box 3. Case study: key factors in successfully implementing TIA

The experience gained by TEWW in implementing the Clinical Link Pathway for Trauma has generated some insights. It has taken a lot of planning, patience and determination to keep it on the agenda in spite of organisational changes, mergers and competing priorities. There are, however, a few key factors that have facilitated the ambitious scope and success so far: first, it was important to sell the concept to senior leaders in the organisation using language that connected with its change processes and aims. The TIA was then sponsored by the medical director. TEWW uses “Lean” methodology, which looks for ways to reduce inefficiencies in its delivery of care. Unidentified trauma was demonstrated using local statistics and service user stories as one way that a person’s journey could stall, be misguided or be less than optimal. The TIA needed to demonstrate how it fitted with the organisation’s key objectives both strategically and in practise with individual clients. By engaging senior support, the approach has maintained high level support in spite of competing demands. Second, it helped to use the methodology for system change that the Trust already employed. TEWW uses “pathways” to describe the structures, management systems and clinical decision making necessary to support the needs of a specific client group. Pathways aim to deliver care which adds value to the client’s health outcome. Most pathways are diagnostic. However, the Trauma Informed Service pathway describes care for anyone showing the effects of trauma regardless of their diagnosis. Third, change is facilitated when staff are empowered by it rather than burdened. This pathway does not dictate what must be done nor is it strictly governed. The process of becoming trauma-informed has become embedded when it is owned by staff and this has had to be gradual over time as awareness deepened. It helped to have emphasis on flexibility of response, to enhance skills and confidence, and to keep data collection to a minimum. Fourth, it helped the pathway lead to understand theory relating to organisational functioning as well as individual trauma work and consider what leadership behaviours others would follow. Empathic engagement with other staff was needed to create alliances and fit the pathway to their clinical needs. Staff needed to be shown the difference it could make to clients and to their own work. They needed opportunity to develop skills, to embed the value of TIA into their own motivational system and to see TIA as a group that they want to belong to. And finally, the personal voices and experiences of service users have been vital in showing the way.

the causes of mental distress is vital (Harris and Fallot, 2001). Practitioners must move from asking “what is wrong with you” to “what happened to you” (Harris and Fallot, 2001). In other words, practitioners must understand the critical and primary role of trauma and fundamentally change their practice as a result. Without this, TIAs are at high risk of co-optation (as has arguably occurred with the concepts of recovery and peer support), meaning that mainstream implementation could be tokenistic, fragmentary and divorced from TIAs core principles.

Although individual practitioners can engage with people in trauma-informed ways, this will be inadequate without system-wide change as systems, as well as individual managers, can block what is needed to effect change (Kotter and Cohen, 2003). Rose *et al.* (2012) assessed DH documentation for its fit with TIA principles. They found that it is policy that trauma and abuse is discussed and documented in all mental health assessments. The principles of TIA are also consistent with policy advocating partnership working between survivors and providers, such as choice and coproduction. However, further policies are needed so that services can move away from force, coercion and risk-averse practice and towards TIAs, with trusts incentivised to implement change.

AK played a key role in introducing TIAs to her mental health trust, and Box 3 explores some of the key factors that made this successful. As can be seen, it was particularly important to demonstrate that TIAs fitted with and furthered key organisational objectives.

Alongside a paradigm shift, there must be discussion and acknowledgement of the critical roles of historical and cultural violence, including ethnically and socio-demographically based differential experiences within the mental health system (e.g. Morgan *et al.*, 2004; McKenzie and Bhui, 2007).

We must combat the view that TIAs are utopian because survivors are dangerous and in need of compulsion (e.g. Muskett, 2014). We must recognise the cycle of crisis that a focus on risk management perpetuates as people struggle for personal agency, choice and control over their lives (Samele *et al.*, 2007). In TEWW, positive risk-taking is policy, including understanding some risky behaviours as survival strategies.

TIA is a complex concept, and opportunities to acquire and develop knowledge are needed. The two 2014 UK conferences on trauma-informed mental healthcare have helped create such opportunities. Yet knowledge alone is insufficient: as Kotter and Cohen (2003) argue, “Without conviction that you can make change happen, you will not act, even if you see the vision”. Support for implementation is also crucial. It is estimated that it takes 10-15 years for new healthcare innovations to be incorporated into routine clinical practice (e.g. Proctor *et al.*, 2009). This gap between research and its implementation is referred to as the translational gap (Tansella and Thornicroft, 2009). Implementation science aims to bridge this translational divide. Thus, in bringing TIA to the UK, implementation science will have an important role to play in supporting individuals and organisations to enact change. Whilst implementation science is beyond the scope of this paper to describe (see, e.g. Damschroder *et al.*, 2009), key ingredients might include implementing a rewards and recognition scheme for staff (Kotter and Cohen, 2003; Tansella and Thornicroft, 2009); understanding current organisational culture and the shifts needed to achieve change (Damschroder *et al.*, 2009); and providing case studies of successful implementation to combat hopelessness and bolster confidence that change is possible (Kotter and Cohen, 2003). Specific to TIA, Harris and Fallot (2009) have developed a self-assessment and planning protocol which supports an organisation’s implementation of TIAs. They argue that if the principles of TIA are reflected in the culture of an organisation for each contact, physical setting, relationship and activity for survivors and staff, the organisation is trauma-informed. Several key steps for moving towards this culture are described in Table III, and Table IV contains a summary of useful resources. In Box 2, AK described her experiences of bringing TIA to TEWW. Key steps in implementing TIA in TEWW have included designing a trauma-informed pathway, training staff, conducting evaluations, developing written guidelines for stakeholders and promoting ownership at senior levels; this implementation can take ten years (Brooker *et al.*, 2016).

Once a critical mass develops, it will become easier for people to model TIA, mentor others, create networks, identify trauma champions and share ideas (Turner, 1990). We believe

Table III Harris and Fallo's key steps for implementing TIA in an organisation

Key step	Activity
Planning	Including leadership commitment, formation of a trauma workgroup to lead and oversee change and the identification of a trauma champion
An initial training event	For as many staff as possible plus service users, encompassing the principles and practice of TIA, care and support for staff, trauma work in the organisation, future directions and implementation
Short-term follow up	The trauma workgroup develops an implementation plan using the protocol, and further training is provided to staff
Longer term follow up	Progress is reviewed including barriers to implementation. Ongoing processes are implemented such as TIA questions in service user experience surveys, and implementation plans added to quality assurance processes

Table IV Further resources

Organisation/study	About	Web link
The ACE study	Website for the landmark study into the link between adverse childhood experiences and adult health. Includes links to publications and study materials	www.acestudy.org/
National Center for Trauma Informed Care	America's National Center for promoting and supporting TIC/TIAs. Includes resources, tools, newsletters and more	www.nasmhpd.org/TA/nctic.aspx
Community Connections	Not-for-profit mental health service provider in Washington, US, headed by Roger Fallo and Maxine Harris, the originators of TIAs. Includes a list of Community Connections products	www.communityconnectionsdc.org
The Anna Institute	Created by Ann Jennings, PhD, whose daughter died by suicide following childhood sexual abuse and enforced detention and treatment in the psychiatric system. Includes resources and links to publications. Ann Jennings hosts a discussion list for people interested in TIAs (SPSCOT)	www.theannainstitute.org
National Council for Community Behavioral Healthcare	As well as having a section for TIAs on their website which lists resources, an issue of the <i>National Council Magazine</i> (2011, issue 2) is dedicated to TIAs	www.thenationalcouncil.org/areas-of-expertise/trauma-informed-behavioral-healthcare/ ; www.thenationalcouncil.org/wp-content/uploads/2012/11/NC-Mag-Trauma-Web-Email.pdf

that trauma survivors have a pivotal role to play in this. We live the impact of trauma everyday. We understand its devastating effects, the damage inflicted by the current mental health system, the need for mutual relationships based on safety and cooperation, the need for personal control, and the vital support of peers. In bringing TIAs to the UK, we need survivor leaders and champions advocating for values-based system change with passion and commitment. Our hope is that this vision will become a reality and that this discussion paper will have contributed to this.

Acknowledgement

Angela Sweeney is funded by a Post-Doctoral Research Fellowship award from the National Institute for Health Research. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health. Dr Sweeney would like to thank the Advisory Groups for their contribution to the APTT study (understanding and improving Assessment Processes for Talking Therapies) from which this paper has arisen: Vanessa Anenden, Katie Bogart, Dr Sarah Carr, Dr Jocelyn Catty, Professor David Clark, Dr Sarah Clement (co-author), Alison Faulkner, Sarah Gibson, Mary Ion, Dr Jayasree Kalathil, Steve Keeble, Dr Angela Kennedy (co-author), Dr Gemma Kothari and Lana Samuels.

References

- Anda, R.F., Butchart, A., Felitti, V.J. and Brown, D.W. (2010), "Building a framework for global surveillance of the public health: implications of adverse childhood experiences", *Preventive Medicine*, Vol. 39 No. 1, pp. 93-8.
- Angermeyer, M.C. and Dietrich, S. (2006), "Public beliefs about and attitudes towards people with mental illness: a review of population studies", *Acta Psychiatrica Scandinavica*, Vol. 113 No. 3, pp. 163-79.
- Ardino, V. (2014), "Trauma-informed care: is cultural competence a viable solution for efficient policy strategies?", *Clinical Neuropsychiatry*, Vol. 11 No. 1, pp. 45-51.
- Azeem, M.W., Aujla, A., Rammerth, M., Binsfeld, G. and Jones, R.B. (2011), "Effectiveness of six core strategies based on trauma informed care in reducing seclusions and restraints at a child and adolescent psychiatric hospital", *Journal of Child & Adolescent Psychiatric Nursing*, Vol. 24 No. 1, pp. 11-5.
- Bentall, R., de Sousa, P., Varese, F., Wickham, S., Sitko, K., Haarmans, M. and Read, J. (2014), "From adversity to psychosis: pathways and mechanisms from specific adversities to specific symptoms", *Social Psychiatry and Psychiatric Epidemiology*, Vol. 49 No. 7, pp. 1011-22.
- Blanch, A., Filson, B., Penney, D. and Cave, C. (2012), *Engaging Women in Trauma-informed Peer Support: A Guidebook*, National Center for Trauma-Informed Care, Rockville, MD.
- Bloom, S. (2006), "Human service systems and organizational stress: thinking and feeling our way out of existing organizational dilemmas", Report for the Trauma Task Force, Philadelphia, PA, available at: www.sanctuaryweb.com/Portals/0/Bloom%20Pubs/2006%20Bloom%20Human%20Service%20Systems%20and%20Organizational%20Stress.pdf (accessed 17 August 2016)
- Bloom, S. and Farragher, B. (2010), *Destroying Sanctuary: The Crisis in Human Service Delivery Systems*, Oxford University Press, New York, NY.
- Bloom, S.L. (2013), *Creating Sanctuary: Toward the Evolution of Sane Societies*, rev. ed., Routledge, New York, NY.
- Bloom, S.L. and Norton, K. (2004), "Special section on the therapeutic community in the 21st century", *Psychiatric Quarterly*, Vol. 75 No. 3, pp. 229-31.
- Brooker, C., Brown, M., Kennedy, A. and Tocque, K. (2016), "The care programme approach, sexual violence and clinical practice in mental health", *Journal of Forensic and Legal Medicine*, Vol. 43, October, pp. 97-101.
- Chambers, M., Gallagher, A., Borschmann, R., Gillard, S., Turner, K. and Kantaris, X. (2014), "The experiences of detained mental health service users: issues of dignity in care", *BMC Medical Ethics*, Vol. 15 No. 50, pp. 1-8, available at: www.biomedcentral.com/content/pdf/1472-6939-15-50.pdf
- Chandler, G. (2008), "From traditional inpatient to trauma-informed treatment: transferring control from staff to patient", *Journal of the American Psychiatric Nurses Association*, Vol. 14 No. 5, pp. 363-71.
- Coles, S. (2014), "Facing the unspeakable", paper presented at the Trauma Informed Services: The Future for Mental Health? Conference, Nottingham, 5 November.
- Cole, S., Eisner, A., Gregory, M. and Ristuccia, J. (2013), *Helping Traumatized Children Learn 2: Creating and Advocating for Trauma-sensitive Schools*, Massachusetts Advocates for Children, Boston, MA.
- Damschroder, L., Aron, D., Keith, R., Kirsh, S., Alexander, J. and Lowery, J. (2009), "Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science", *Implementation Science*, Vol. 4 No. 50, doi: 10.1186/1748-5908-4-50.
- DH (2003), *Mainstreaming Gender and Women's Mental Health: Implementation Guidance*, Department of Health, London.
- DH (2011), *Commissioning Services for Women and Children who are Victims of Violence or Abuse – A Guide for Health Commissioners*, Department of Health, London.
- Dillon, J., Johnstone, L. and Longden, E. (2012), "Trauma, dissociation, attachment & neuroscience: a new paradigm for understanding severe mental distress", *The Journal of Critical Psychology, Counselling and Psychotherapy*, Vol. 12 No. 3, pp. 145-55.
- Dolezal, T., McCollum, D., Callahan, M. and Eden, P. (2009), "Hidden costs in health care: the economic impact of violence and abuse", Academy on Violence and Abuse, MN.

- Domino, M.E., Morrissey, J.P., Chung, S. and Nadlicki, T. (2006), "Changes in service use during a trauma-informed intervention for women", *Women and Health*, Vol. 44 No. 3, pp. 105-22.
- Durant, J.J. (2011), "Dissociation and its discontents: an exploration of the role of dissociation in a retraumatization cycle among victims of repeated trauma", doctoral dissertation, California Institute of Integral Studies, San Francisco, CA.
- Elliot, D., Bjelajac, P., Fallot, R., Markoff, L. and Glover Reed, B. (2005), "Trauma-informed or trauma-denied: principles and implementation of trauma-informed services for women", *Journal of Community Psychology*, Vol. 33 No. 4, pp. 461-77.
- Esaki, N. and Larkin, H. (2013), "Prevalence of adverse childhood experiences (ACEs) among child service providers", *Families in Society: The Journal of Contemporary Social Services*, Vol. 94 No. 1, pp. 31-7.
- Filson, B. (2011), "Is anyone really listening?", *National Council Magazine. Special Issue: Breaking the Silence: Trauma-informed Behavioral Healthcare*, Nos 2/15, p. 15.
- Filson, B. and Mead, S. (2016), "Intentional peer support: an alternative approach", in Russo, J. and Sweeney, A. (Eds), *Searching for a Rose Garden: Fostering real Alternatives to Psychiatry*, PCCS Books, Ross-on-Wye, pp. 109-17.
- Fisher, H., Jones, P.B., Fearon, P., Craig, T., Dazzan, P., Morgan, K., Hutchinson, G., Doody, G.A., McGuffin, P., Leff, J., Murry, R.M. and Morgan, C. (2010), "The varying impact of type, timing and frequency of exposure to childhood adversity on its association with adult psychotic disorder", *Psychological Medicine*, Vol. 40 No. 12, pp. 1967-78.
- Freuh, C., Knapp, R., Cusack, K., Grubaugh, A., Sauvageot, J., Cousins, V., Yim, E., Robins, C., Monnier, J. and Hiers, T. (2005), "Patients' reports of traumatic or harmful experiences within the psychiatric setting", *Psychiatric Services*, Vol. 56 No. 9, pp. 1123-33.
- Friedman, M. (2014), "Gay conversion therapy: a dark chapter in mental health care", *Psychology Today*, 21 January, available at: www.psychologytoday.com/blog/brick-brick/201401/gay-conversion-therapy-dark-chapter-in-mental-health-care
- Gatz, M., Brown, V., Hennigan, K., Rechberger, E., O'Keefe, M., Rose, T. and Bjelajac, P. (2007), "Effectiveness of an integrated, trauma-informed approach to treating women with co-occurring disorders and histories of trauma: the Los Angeles site experience", *Journal of Community Psychology*, Vol. 35 No. 7, pp. 863-78.
- Greenwald, R., Siradas, L., Schmitt, T., Reslan, S., Fierle, J. and Sande, B. (2012), "Implementing trauma-informed treatment for youth in a residential facility: first-year outcomes", *Residential Treatment for Children and Youth*, Vol. 29 No. 2, pp. 141-53.
- Harragan, S. (2013), "Therapeutic residential care for children and young people: an attachment and trauma-informed model for practice", *Adoption and Fostering*, Vol. 37 No. 1, pp. 102-3.
- Harris, M. and Fallot, R. (2001), *Using Trauma Theory to Design Service Systems. New Directions for Mental Health Services*, Jossey-Bass, San Francisco, CA.
- Harris, R. and Fallot, M. (2009), *Creating Cultures of Trauma-informed Care (CCTIC): A Self-Assessment and Planning Protocol*, Community Connections, Washington, DC.
- Hatch, S. and Dohrenwend, B. (2007), "Distribution of traumatic and other stressful life events by race/ethnicity, gender, SES and age: a review of the research", *American Journal of Community Psychology*, Vol. 40 Nos 3-4, pp. 313-32.
- Havig, K. (2008), "The health care experiences of adult survivors of child sexual abuse: a systematic review of evidence on sensitive practice", *Trauma, Violence, & Abuse*, Vol. 9 No. 1, pp. 19-33.
- Hepworth, I. and McGowan, L. (2013), "Do mental health professionals enquire about childhood sexual abuse during routine mental health assessment in acute mental health settings? A substantive literature review", *Journal of Psychiatric and Mental Health Nursing*, Vol. 20 No. 6, pp. 473-83.
- IIMHL (2012), *Make It So: Key National Activities in Trauma-informed Care across IIMHL Countries*, International Initiative for Mental Health Leadership, Pakuranga, Auckland.
- Jackson, V. (2003), "In our own voice: African-American stories of oppression, survival and recovery in mental health systems", *Off Our Backs*, Vol. 33 Nos 7/8, pp. 19-21.

- Kessler, R., McLaughlin, K., Green, J., Gruber, M.J., Sampson, N.A., Zaslavsky, A.M., Aguilar-Gaxiola, S., Alhamzawi, A.O., Alonso, J., Angermeyer, M., Benjet, C., Bromet, E., Chatterji, S., de Girolamo, G., Demyttenaere, K., Fayyad, J., Florescu, S., Gal, G., Gureje, O., Haro, J.M., Hu, C.Y., Karam, E.G., Kawakami, N., Lee, S., Lépine, J.P., Ormel, J., Posada-Villa, J., Sagar, R., Tsang, A., Ustün, T.B., Vassilev, S., Viana, M.C. and Williams, D.R. (2010), "Childhood adversities and adult psychopathology in the WHO world mental health surveys", *British Journal of Psychiatry*, Vol. 197 No. 5, pp. 378-85.
- Khalifeh, H., Moran, P., Borschmann, R., Dean, K., Hart, C., Hogg, J., Osborn, O., Johnson, S. and Howard, L. (2015), "Domestic and sexual violence against patients with severe mental illness", *Psychological Medicine*, Vol. 45 No. 4, pp. 875-86.
- Knight, C. (2015), "Trauma-informed social work practice: practice considerations and challenges", *Clinical Social Work Journal*, Vol. 43, pp. 25-37.
- Kotter, J. and Cohen, D. (2003), "Creative ways to empower action to change the organization: cases in point", *Journal of Organizational Excellence*, Vol. 22 No. 2, pp. 101-10.
- LeBel, J. and Goldstein, R. (2006), "The economic cost of using restraint and the value added by restraint reduction or elimination", *Psychiatric Services*, Vol. 56 No. 9, pp. 1109-14.
- McKenzie, K. and Bhui, K. (2007), "Institutional racism in mental health care", *British Medical Journal*, Vol. 334 No. 7595, pp. 649-50.
- McNeish, D. and Scott, S. (2008), "Meeting the needs of survivors of abuse: Mental Health Trusts Collaboration Project overview of evaluation findings", Department of Health/National Institute of Mental Health, London, available at: www.dmss.co.uk/wp-content/uploads/2013/06/Final-overview-report.pdf (accessed 17 August 2016)
- Mauritz, M., Goossens, P., Draijer, N. and van Achterberg, T. (2013), "Prevalence of interpersonal trauma exposure and trauma-related disorders in severe mental illness", *European Journal of Psychotraumatology*, Vol. 4, available at: www.ncbi.nlm.nih.gov/pmc/articles/PMC3621904/
- Mead, S. and MacNeil, C. (2006), "Peer support: what makes it unique?", *International Journal of Psychosocial Rehabilitation*, Vol. 10 No. 2, pp. 29-37.
- Messina, N., Calhoun, S. and Braithwaite, J. (2014), "Trauma-informed treatment decreases posttraumatic stress disorder among women offenders", *Journal of Trauma and Dissociation*, Vol. 15 No. 1, pp. 6-23.
- Mohan, R., McCrone, P., Szmukler, G., Micali, N., Afuwape, S. and Thornicroft, G. (2006), "Ethnic differences in mental health service use among patients with psychotic disorders", *Social Psychiatry and Psychiatric Epidemiology*, Vol. 41 No. 10, pp. 771-6.
- Morgan, C., Mallett, R., Hutchinson, G. and Leff, J. (2004), "Negative pathways to psychiatric care and ethnicity: the bridge between social science and psychiatry", *Social Science and Medicine*, Vol. 58 No. 4, pp. 739-52.
- Morrissey, J., Jackson, E., Ellis, A., Amaro, H., Brown, V. and Najavits, L. (2005), "Twelve-month outcomes of trauma-informed interventions for women with co-occurring disorders", *Psychiatric Services*, Vol. 56 No. 10, pp. 1213-22.
- Moskowitz, A. (2011), "Schizophrenia, trauma, dissociation, and scientific revolutions", *Journal of Trauma and Dissociation*, Vol. 12 No. 4, pp. 347-57.
- Muskett, C. (2014), "Trauma-informed care in inpatient mental health settings: a review of the literature", *International Journal of Mental Health Nursing*, Vol. 23 No. 1, pp. 51-9.
- National Mental Health Development Unit (2010), *Working towards Women's Well-being: Unfinished Business*, National Mental Health Development Unit, London.
- NICE (2005), *Violence: the Short-Term Management of Disturbed/Violent Behaviour in In-Patient Psychiatric Settings and Emergency Departments*, Guidance CG25, National Institute for Health and Care Excellence, London.
- NICE (2014), *Psychosis and Schizophrenia in Adults: Treatment and Management*, Guidance CG178, National Institute for Health and Care Excellence, London.
- O'Hagan, M. (2003), "Force in mental health services: international user/survivor perspectives", Keynote Address to the World Federation for Mental Health Biennial Congress, Melbourne.
- Paradies, Y. (2006), "A systematic review of empirical research on self-reported racism and health", *International Journal of Epidemiology*, Vol. 35 No. 4, pp. 888-901.

- Paterson, B. (2014), "Mainstreaming trauma", paper presented at the Psychological Trauma-Informed Care Conference, Stirling University, Stirling, 4 June, available at: www.stir.ac.uk/media/schools/nursing/documents/Trauma14-Paterson-mainstreaming-trauma-workshop.pdf
- Paterson, B., McIntosh, I., Wilkinson, D., McComish, S. and Smith, I. (2013), "Corrupted cultures in mental health inpatient settings. Is restraint reduction the answer?", *Journal of Psychiatric Mental Health Nursing*, Vol. 20 No. 3, pp. 228-35.
- Perry, B. (2005), "Maltreatment and the developing child: how early childhood experience shapes child and culture", The Inaugural Margaret McCain lecture (abstracted), McCain Lecture Series, The Centre for Children and Families in the Justice System, London.
- Proctor, E., Landsverk, J., Aarons, G., Chambers, D., Glisson, C. and Mittman, B. (2009), "Implementation research in mental health services: an emerging science with conceptual, methodological, and training challenges", *Administration and Policy in Mental Health*, Vol. 36 No. 1, pp. 24-34.
- Read, J. (2010), "Can poverty drive you mad? 'Schizophrenia', socio-economic status and the case for primary prevention", *New Zealand Journal of Psychology*, Vol. 39 No. 2, pp. 7-19.
- Read, J., Bentall, R. and Fosse, R. (2009), "Time to abandon the bio-bio-bio model of psychosis: exploring the epigenetic and psychological mechanisms by which adverse life events lead to psychotic symptoms", *Epidemiologia e Psichiatria Sociale*, Vol. 18 No. 4, pp. 299-310.
- Read, J., Hammersley, P. and Rudegeair, T. (2007), "Why, when and how to ask about childhood abuse", *Advances in Psychiatric Treatment*, Vol. 13, pp. 101-10.
- Read, J., Magliano, L. and Beavan, V. (2013), "Public beliefs about the causes of 'schizophrenia': bad things happen and can drive you crazy", in Read, J. and Bentall, R. (Eds), *Models of Madness: Psychological, Social and Biological Approaches to Schizophrenia*, 2nd ed., Brunner-Routledge, Hove and New York, NY, pp. 133-46.
- Read, J., Fosse, R., Moskowitz, A. and Perry, B. (2014), "The traumagenic neurodevelopmental model of psychosis revisited", *Neuropsychiatry*, Vol. 4 No. 1, pp. 65-79.
- Rose, S., Freeman, C. and Proudlock, S. (2012), "Despite the evidence – why are we still not creating more trauma informed mental health services?", *Journal of Public Mental Health*, Vol. 11 No. 1, pp. 5-9.
- Samele, C., Lawton-Smith, S., Warner, L. and Mariathasan, J. (2007), "Patient choice in psychiatry", *The British Journal of Psychiatry*, Vol. 191 No. 1, pp. 1-2.
- SAMHSA (2014), *SAMHSA's Working Concept of Trauma and Framework for a Trauma-Informed Approach*, National Centre for Trauma-Informed Care (NCTIC), SAMHSA, Rockville, MD.
- Schachter, C.L., Stalker, C.A., Teram, E., Lasiuk, G.C. and Danilkewich, A. (2008), *Handbook on Sensitive Practice for Health Care Practitioner: Lessons from Adult Survivors of Childhood Sexual Abuse*, Public Health Agency of Canada, Ottawa.
- Scottish Government (2012), "Scotland's mental health strategy 2012-2015", Scottish Government, Edinburgh, available at: www.scotland.gov.uk/Resource/0039/00398762.pdf (accessed 17 August 2016).
- Shanock, L.R. and Eisenberger, R. (2006), "When supervisors feel supported: relationships with subordinates' perceived supervisor support, perceived organizational support, and performance", *Journal of Applied Psychology*, Vol. 91 No. 3, pp. 689-95.
- Shevlin, M., Houston, J.E., Dorahy, M.J. and Adamson, G. (2008), "Cumulative traumas and psychosis: an analysis of the national comorbidity survey and the British psychiatric morbidity survey", *Schizophrenia Bulletin*, Vol. 34 No. 1, pp. 193-9.
- St-Amand, N. and LeBlanc, E. (2013), "Women in 19th century asylums: three exemplary women; a new Brunswick hero", in Le Francois, B., Menzies, R. and Reaume, G. (Eds), *Mad Matters: A Critical Reader in Canadian Mad Studies*, Canadian Scholars' Press Inc., Toronto, pp. 38-48.
- Steckley, L. (2013), "Therapeutic residential child care for children and young people: an attachment and trauma-informed model for practice", *British Journal of Social Work*, Vol. 43 No. 1, pp. 199-201.
- Tansella, M. and Thornicroft, G. (2009), "Implementation science: understanding the translation of evidence into practice", *British Journal of Psychiatry*, Vol. 195 No. 4, pp. 283-5.

Taylor, C. (2012), *Empathic Care for Children with Disorganized Attachments: a Model for Mentalizing, Attachment and Trauma-informed Care*, Jessica Kingsley, London.

Terr, L. (1991), "Childhood traumas: an outline and overview", *American Journal of Psychiatry*, Vol. 148 No. 1, pp. 10-20.

Toner, J., Daiches, A. and Larkin, W. (2013), "Asking about trauma: the experiences of psychological therapists in early intervention services", *Psychosis*, Vol. 5 No. 2, pp. 175-86.

Turner, R. (1990), "Role change", *Annual Review of Sociology*, Vol. 16 No. 1, pp. 87-110.

Van der Kolk, B.A. (2005), "Developmental trauma disorder: towards a rational diagnosis for chronically traumatized children", *Psychiatric Annals*, Vol. 35 No. 5, pp. 401-8.

Varese, F., Smeets, F., Drukkers, M., Lieveerse, R., Lataster, T., Viechtbauer, W., Read, J., van Os, J. and Bentall, R. (2012), "Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective and cross-sectional cohort studies", *Schizophrenia Bulletin*, Vol. 38 No. 4, pp. 661-71.

Wardhaugh, J. and Wilding, P. (1993), "Towards an explanation of the corruption of care", *Critical Social Policy*, Vol. 13 No. 1, pp. 4-32.

Weissbecker, I. and Clark, C. (2007), "The impact of violence and abuse on women's physical health: can trauma-informed treatment make a difference?", *Journal of Community Psychology*, Vol. 35 No. 7, pp. 909-23.

Wilson, A. and Beresford, P. (2002), "Madness, distress and postmodernity: putting the record straight", in Corker, M. and Shakespeare, T. (Eds), *Disability/Postmodernity: Embodying Disability Theory*, Continuum, New York NY, pp. 143-58.

Web references

www.nctsn.net/trauma-types/complex-trauma

www.cdc.gov/violenceprevention/acestudy/prevalence.html

www.theannainstitute.com

www.newfocas.co.uk/includes/spaw2/uploads/files/trauma.pdf

www.beyondyouthcustody.net/wp-content/uploads/BYC-Developing-trauma-resettlement-youth-custody-practitioners-guide.pdf

<http://stephaniecovington.com/assets/files/Brochure-Flyer-dated-2-26-14-Woman-at-Risk-Flyer-V1.pdf>

www.vptorg.co.uk/whoslistening/docs/conf%20speeches/Aimee_Neri_Presentation.pdf

www.evoc.org.uk/wordpress/wp-content/media/2014/07/Jenney-seminar.pdf

www.makingwaves.org/news/trauma-informed-services-conference/

www.stir.ac.uk/trauma14/

www.bild.org.uk/our-services/events/supportingpeople/

www.ukpts.co.uk/site/assets/CALM-Final.pdf

www.calmtraining.co.uk/index.php

www.ewrasac.org.uk/Training/

www.avaproject.org.uk

www.starwards.org.uk

www.safewards.net/

Further reading

Radmore, J. (2013), "The effect of initiating trauma informed care on clients' perception of safety", *Masters Abstracts International*, Vol. 51 No. 4, pp. 1-49.

Author affiliations

Angela Sweeney is a NIHR Post-Doctoral Research Fellow at Population Health Research Institute, St George's University of London, London, UK.

Sarah Clement is Visiting Researcher at the Department of Health Service and Population Research, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK.

Beth Filson is an Independent Writer and Trainer in Trauma-informed Approaches, Williamsburg, Massachusetts, USA.

Angela Kennedy is a Consultant Clinical Psychologist and Service Lead at Tertiary Psychosis Service, Tees, Esk and Wear Valleys NHS Trust, Durham, UK.

Corresponding author

Angela Sweeney can be contacted at: asweeney@sgul.ac.uk

For instructions on how to order reprints of this article, please visit our website:

www.emeraldgroupublishing.com/licensing/reprints.htm

Or contact us for further details: permissions@emeraldinsight.com

This article has been cited by:

1. John Read, David Harper, Ian Tucker, Angela Kennedy. 2018. How do mental health services respond when child abuse or neglect become known? A literature review. *International Journal of Mental Health Nursing* 11. . [[Crossref](#)]
2. Pauline Cusack, Frank Patrick Cusack, Sue McAndrew, Mick McKeown, Joy Duxbury. 2018. An integrative review exploring the physical and psychological harm inherent in using restraint in mental health inpatient settings. *International Journal of Mental Health Nursing* 27:3, 1162-1176. [[Crossref](#)]
3. Ian Cummins. 2018. The Impact of Austerity on Mental Health Service Provision: A UK Perspective. *International Journal of Environmental Research and Public Health* 15:6, 1145. [[Crossref](#)]
4. Nicola Atwool. 2018. Challenges of operationalizing trauma-informed practice in child protection services in New Zealand. *Child & Family Social Work* 40. . [[Crossref](#)]
5. Nicky Lambert. 2018. Supporting women with mental health issues. *Mental Health Practice* . [[Crossref](#)]
6. FoleyMarian, Marian Foley, CumminsIan, Ian Cummins. 2018. Reporting sexual violence on mental health wards. *The Journal of Adult Protection* 20:2, 93-100. [[Abstract](#)] [[Full Text](#)] [[PDF](#)]
7. Diana Rose, Sarah Carr, Peter Beresford. 2018. 'Widening cross-disciplinary research for mental health': what is missing from the Research Councils UK mental health agenda?. *Disability & Society* 33:3, 476-481. [[Crossref](#)]
8. John Read. 2018. Making Sense of, and Responding Sensibly to, Psychosis. *Journal of Humanistic Psychology* 9, 002216781876191. [[Crossref](#)]
9. John Read, David Harper, Ian Tucker, Angela Kennedy. 2018. Do adult mental health services identify child abuse and neglect? A systematic review. *International Journal of Mental Health Nursing* 27:1, 7-19. [[Crossref](#)]
10. Edward J. Alessi, Sarilee Kahn. 2017. Using psychodynamic interventions to engage in trauma-informed practice. *Journal of Social Work Practice* 31, 1-13. [[Crossref](#)]
11. Sarah Carr. 2017. Renegotiating the contract. *The Lancet Psychiatry* 4:10, 740-741. [[Crossref](#)]
12. Vittoria Ardino. 2017. Healthcare utilization and trauma-informed care: A discussion paper. *MALTRATTAMENTO E ABUSO ALL'INFANZIA* :2, 31-53. [[Crossref](#)]
13. Samantha McEvedy, Tessa Maguire, Trentham Furness, Brian McKenna. 2017. Sensory modulation and trauma-informed-care knowledge transfer and translation in mental health services in Victoria: Evaluation of a statewide train-the-trainer intervention. *Nurse Education in Practice* 25, 36-42. [[Crossref](#)]
14. SpandlerHelen, Helen Spandler, McKeownMick, Mick Mckeown. 2017. Exploring the case for truth and reconciliation in mental health services. *Mental Health Review Journal* 22:2, 83-94. [[Abstract](#)] [[Full Text](#)] [[PDF](#)]